

Bose-Einstein condensation in systems with quadratic energy levels

José M. B. Noronha* and David J. Toms†
*Department of Physics, University of Newcastle upon Tyne,
 Newcastle upon Tyne, NE1 7RU, U. K.
 (June 13, 2001)*

We study finite size effects on ideal Bose gas thermodynamics (and where applicable Bose-Einstein condensation) in several systems with quadratic particle energy levels, namely, the Einstein universe, the one-dimensional box and the three-dimensional flat box ($1\text{Dbox} \otimes \mathbb{R}^2$). The Mellin-Barnes transform technique is used to obtain analytical approximations to thermodynamic quantities, providing an analytical description of the specific heat peak and the onset of condensation.

05.30.Jp, 03.75.Fi

I. INTRODUCTION

Since the experimental realization of Bose-Einstein condensation (BEC) for a dilute atomic gas in 1995, this field has been the subject of intensive research, both theoretical and experimental. (See [1] and [2] for theoretical and experimental reviews respectively.)

In particular, finite size effects have been dealt with in a number of works, both numerically and using a variety of analytical techniques. Rigorously speaking, the critical behaviour known as BEC may only happen when we take the limit $N \rightarrow \infty$ (N being the number of particles). For a finite system all thermodynamic quantities are analytical. As the size of the system is taken to infinity in some well defined way (e.g., fixed temperature and density in the case of a homogeneous system), it is possible that the chemical potential (seen as a function of the temperature T when N is given) will approach a non-analytical function at some critical temperature T_c . This will cause other thermodynamic quantities to also approach non-analytical behaviour, which can result in a phase transition.

In this context, the harmonic oscillator trapping potential has received special attention [3–7], motivated by its suitability to describe experimental traps and the simple form of the energy eigenvalues. The three-sphere S^3 (spatial section of the Einstein universe) has also received some attention. In this case the form of the energy eigenvalues is still simple, but the fact that it involves squares of the quantum numbers describing the state makes the problem of treating the thermodynamic sums more complicated. In fact, it is extremely complicated if the gas is considered to be relativistic. There is not to this date a study of the relativistic Bose gas in S^3 relying mostly on analytical expressions. Altaie [8] was the first to provide an analytical study of the non-relativistic case. Singh and Pathria [9] studied the relativistic case obtaining Altaie's result in the non-relativistic limit. More recently, Trucks [10] repeated the calculations of [9] for the minimally coupled case (whereas [9] had done it for conformal coupling) and Altaie and Malkawi [11] for the spin-1 case (whereas [9] had done it for spin-0). Other authors have studied the problem of a Bose gas in an Einstein universe (and more generally, curved spaces) from different perspectives, not focusing on finite size effects [12–15].

Finite size effects for bosons trapped in a potential (including boxes) have also been studied analytically in a general setting [16]. We will discuss this setting in more detail later. The basic idea behind the work of [16], the Mellin-Barnes transform, is common to our work as well. There is nevertheless an important difference that makes the method of [16] not applicable to the situations concerned here.

All the investigations we have mentioned considered the particles to be non-interacting. Finite size effects have also been taken into account for bosons with self-interactions, but only numerically [17].

In this article, we make use of the Mellin-Barnes transform (used in similar contexts in the past in [6,16,18,19]) to study the thermodynamics of an ideal non-relativistic spin-0 Bose gas in three systems that present some common features in the form of the energy eigenvalues: they are all quadratic. These systems are the three-sphere (non-relativistic spin-0 gas), the one-dimensional box (1Dbox) and the $1\text{Dbox} \otimes \mathbb{R}^2$. (By this last example we mean a 3-dimensional box, which is finite in one spatial direction and infinite in the other two.)

*E-mail: josenoronha@hotmail.com

†d.j.toms@ncl.ac.uk

In section II, we use the Mellin-Barnes transform to obtain expressions for the number of particles and specific heat. In section III, we briefly discuss some matters inherent to this technique. In section IV, we expand the results of section II around the critical region to obtain the chemical potential as an explicit function of the temperature. From this, we obtain the specific heat as a function of the temperature only and consequently, we have an analytical description of the sharpening of the specific heat peak. In section V are some concluding remarks and a summary of the most important points in the article.

II. NUMBER OF PARTICLES AND SPECIFIC HEAT

Consider a system of non-self-interacting bosons with particle eigenstates of energy E_n . For a relatively high number of particles, and if we are not interested in fluctuations, which is the case, the thermodynamics of this system can be well described using grand-canonical statistics [19–24], greatly simplifying matters. The number of particles is then

$$N = \sum_n [e^{\beta(E_n - \mu)} - 1]^{-1}, \quad (1)$$

where β is the inverse of the temperature T , μ is the chemical potential and the sum is over all particle eigenstates. This expression gives us μ implicitly if we fix T and N . In this section, we shall aim at obtaining analytical expressions for N and the specific heat, C , as functions of μ and β . Rather than presenting the method in a general setting we prefer to develop it first for one specific case, the Einstein universe, and then generalize it to our other applications. We will use the natural units system ($\hbar = 1$ and $c = 1$) and $k_B = 1$ throughout (k_B being Boltzmann's constant).

A. The Einstein Universe

In this case the energy eigenvalues are given by

$$E_n = \frac{1}{a} [n^2 + m^2 a^2 + 6\xi - 1]^{\frac{1}{2}}, \quad n = 1, 2, 3, \dots \quad (2)$$

with degeneracy $d_n = n^2$. a is the radius of the universe, m is the particle mass and ξ is the coupling constant. We will limit ourselves to conformal coupling, $\xi = 1/6$. As noted in [11], this is not very restrictive since choosing a different coupling is equivalent to changing the mass from m to $[m^2 + (6\xi - 1)/a^2]^{1/2}$. For large a , this change is very small. In particular, in the large volume limit, all couplings will give the same results. In the non-relativistic limit we have

$$E_n \simeq \frac{n^2}{2ma^2}. \quad (3)$$

Taking this limit requires $n^2 \ll m^2 a^2$, ie, little occupation of high energy states (“high” when compared with m), corresponding to $T \ll m$. For this to be true in the temperature range we are most interested in, the critical region, it suffices having $\rho \ll m^3$, where ρ is the particle density.

We have then

$$\begin{aligned} N &= \sum_{n=1}^{+\infty} n^2 \left[e^{x(n^2 + \epsilon)} - 1 \right]^{-1} \\ &= \sum_{n=1}^{+\infty} n^2 \sum_{k=1}^{+\infty} e^{-kx(n^2 + \epsilon)}, \end{aligned} \quad (4)$$

where we have changed to the dimensionless variables,

$$x = \frac{\beta}{2ma^2}, \quad \epsilon = -2ma^2\mu. \quad (5)$$

(Note that $\epsilon > -1$ always and $\epsilon \rightarrow -1$ implies $N \rightarrow +\infty$.) At this stage we invoke the Mellin-Barnes transform,

$$e^{-x} = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} d\alpha \Gamma(\alpha) x^{-\alpha}, \quad x > 0, \quad c > 0. \quad (6)$$

Applying it to the exponential of (4) yields

$$N = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} d\alpha \Gamma(\alpha) x^{-\alpha} \sum_{k=1}^{+\infty} k^{-\alpha} \sum_{n=1}^{+\infty} n^2 (n^2 + \epsilon)^{-\alpha}, \quad (7)$$

where we have changed the order of summations and integration. For this we must have $c > 3/2$, so that the sums remain convergent. We now perform the integral by analytically continuing the sums to the whole complex plane (except eventual poles), closing the contour to the left in a large rectangle (corresponding to a high temperature, or small x , expansion) and applying the residue theorem.

$$N \sim \frac{1}{2\pi i} \int_{\gamma} d\alpha \Gamma(\alpha) x^{-\alpha} \zeta(\alpha) f_1(\alpha, \epsilon). \quad (8)$$

γ is the contour mentioned above, $\zeta(\alpha)$ is the Riemann zeta function and $f_i(\alpha, \epsilon)$ is the analytical continuation of the sum $\sum_{n=1}^{+\infty} n^{2i} (n^2 + \epsilon)^{-\alpha}$. The usefulness of defining $f_i(\alpha, \epsilon)$ for general i becomes apparent when we consider other sums later in the article. These functions are treated in the appendix. We use the sign “ $\sim$ ” and not “ $=$ ” as the part of the contour that was added to the integral is not necessarily vanishing. We will come to this later.

In order to extract the poles of the integrand let us focus on $\Gamma(\alpha) f_1(\alpha, \epsilon)$. Consider for now $\alpha > 3/2$. In this case $f_1(\alpha, \epsilon)$ is just the summation we had before. By applying the Γ -function integral representation we have

$$\Gamma(\alpha) f_1(\alpha, \epsilon) = \int_0^{+\infty} dt t^{\alpha-1} e^{-t\epsilon} \sum_{n=1}^{+\infty} n^2 e^{-tn^2}. \quad (9)$$

We can now obtain an expansion for the sum in (9) by using the Mellin-Barnes transform on e^{-tn^2} , yielding

$$\begin{aligned} \sum_{n=1}^{+\infty} n^2 e^{-tn^2} &= \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} d\alpha \Gamma(\alpha) t^{-\alpha} \zeta(2\alpha - 2) \\ &\sim \frac{\sqrt{\pi}}{4} t^{-\frac{3}{2}}, \end{aligned} \quad (10)$$

where we solved the integral by closing the contour to the left and applying the residue theorem. For some properties of the Γ - and Riemann ζ -functions see [25]. Expression (9) now becomes

$$\Gamma(\alpha) f_1(\alpha, \epsilon) \sim \frac{\sqrt{\pi}}{4} \int_0^b dt t^{\alpha-\frac{5}{2}} e^{-t\epsilon} + \frac{\sqrt{\pi}}{4} \int_b^{+\infty} dt t^{\alpha-\frac{5}{2}} e^{-t\epsilon}, \quad (11)$$

where b is a positive number that can be arbitrarily small. The reason we divide the integral in two parts is that since the poles and their residues are the only pieces of information we require, we can concentrate on the first term of (11), which contains them. (The second integral is an analytic function of α .) We will denote the first term by A . Taylor expanding the exponential and performing the integral yields

$$A = \frac{\sqrt{\pi}}{4} \sum_{k=0}^{+\infty} \frac{(-1)^k}{k!} \epsilon^k \frac{b^{\alpha+k-\frac{3}{2}}}{\alpha+k-\frac{3}{2}}. \quad (12)$$

The analytical continuation to the complex plane is now trivial. The poles are at

$$\alpha = \frac{3}{2} - k, \quad k = 0, 1, 2, \dots \quad (13)$$

and the residues at these poles are

$$\text{res}(k) = \frac{\sqrt{\pi}}{4} \frac{(-1)^k}{k!} \epsilon^k. \quad (14)$$

Using this information in (8) results in

$$N \sim \frac{\sqrt{\pi}}{4} x^{-\frac{3}{2}} \sum_{k=0}^{+\infty} \frac{(-1)^k}{k!} \zeta\left(\frac{3}{2} - k\right) (\epsilon x)^k + x^{-1} f_1(1, \epsilon). \quad (15)$$

The sum in this expression comes from the poles of $\Gamma(\alpha)f_1(\alpha, \epsilon)$, and the last term in x^{-1} comes from the pole of $\zeta(\alpha)$. In the high temperature range (small x) it is enough to consider only a few terms from the sum plus the other term. The value of $f_1(1, \epsilon)$ is given in (A14). We finally obtain the approximate result

$$N \sim \frac{\sqrt{\pi}}{4} x^{-\frac{3}{2}} \left[\zeta\left(\frac{3}{2}\right) - \zeta\left(\frac{1}{2}\right)\epsilon x - \frac{1}{4\pi} \zeta\left(\frac{3}{2}\right)(\epsilon x)^2 + \dots \right] - \frac{\pi\sqrt{\epsilon}}{2} \coth(\pi\sqrt{\epsilon})x^{-1}. \quad (16)$$

Instead of having a hyperbolic cotangent in the last term we could have it expressed in terms of a regular cotangent (see appendix). When we take the thermodynamic limit ($V \rightarrow +\infty$, constant density), we have $x \rightarrow 0$ and this result coincides with the continuum approximation, as would be expected.

Altaie [8], using a completely different technique (involving the Poisson summation formula), obtained a result almost identical to ours. The only difference, apart from the total disparity in the method, is that we have additional terms present. However, these new terms could have been found by extending the polylogarithm expansions in Altaie's method to higher order.

We now turn our attention to the specific heat. The total internal energy of the system is given by

$$U = \sum_n E_n \left[e^{\beta(E_n - \mu)} - 1 \right]^{-1}, \quad (17)$$

where the sum is over particle energy eigenstates. In the case of non-relativistic particles in an Einstein universe, it is

$$U = \frac{1}{2ma^2} \sum_{n=1}^{+\infty} n^4 \left[e^{x(n^2 + \epsilon)} - 1 \right]^{-1}. \quad (18)$$

The specific heat at constant radius of the universe is

$$C = \left(\frac{\partial U}{\partial T} \right)_{N,a} = x^2 \left[S(3, 1) - \frac{S(2, 1)^2}{S(1, 1)} \right], \quad (19)$$

where we define a class of sums

$$S(i, j) = \sum_{n=1}^{+\infty} n^{2i} \sum_{k=1}^{+\infty} k^j e^{-kx(n^2 + \epsilon)}. \quad (20)$$

We used $\left(\frac{\partial N}{\partial x} \right)_{N,a} = 0$ to obtain $\left(\frac{\partial(x\epsilon)}{\partial x} \right)_{N,a}$. (For details of how to get (19) see [6].) Applying the Mellin-Barnes transform to the sums $S(i, 1)$ results in

$$S(i, 1) \sim \frac{1}{2\pi i} \int_{\gamma} d\alpha \Gamma(\alpha) x^{-\alpha} \zeta(\alpha - 1) f_i(\alpha, \epsilon), \quad i = 1, 2, 3, \quad (21)$$

as the analogue of (8). Studying $\Gamma(\alpha)f_i(\alpha, \epsilon)$ in the same manner as we did for the case $i = 1$, we find the poles given by

$$\alpha = \frac{1}{2} + i - k, \quad k = 0, 1, 2, \dots \quad (22)$$

The respective residues are

$$\text{res} = \frac{1}{2} \Gamma\left(\frac{2i+1}{2}\right) \frac{(-1)^k}{k!} \epsilon^k. \quad (23)$$

Finally we have the expansion

$$S(i, 1) = \frac{1}{2}\Gamma\left(\frac{2i+1}{2}\right)x^{-\frac{1}{2}-i}\sum_{k=0}^{+\infty}\frac{(-1)^k}{k!}\zeta\left(-\frac{1}{2}+i-k\right)+f_i(2, \epsilon)x^{-2}. \quad (24)$$

Using (A15), (A17) and (A18) for $f_i(2, \epsilon)$ yields

$$\begin{aligned} S(1, 1) &\sim \frac{\sqrt{\pi}}{4}x^{-\frac{3}{2}}\left[\zeta\left(\frac{1}{2}\right)+\frac{1}{4\pi}\zeta\left(\frac{3}{2}\right)x\epsilon-\frac{3}{32\pi^2}\zeta\left(\frac{5}{2}\right)(x\epsilon)^2\right] \\ &\quad +\left[\frac{\pi}{4\sqrt{\epsilon}}\coth(\pi\sqrt{\epsilon})-\frac{\pi^2}{4}\operatorname{csch}^2(\pi\sqrt{\epsilon})\right]x^{-2}, \\ S(2, 1) &\sim \frac{3\sqrt{\pi}}{8}x^{-\frac{5}{2}}\left[\zeta\left(\frac{3}{2}\right)-\zeta\left(\frac{1}{2}\right)x\epsilon-\frac{1}{8\pi}\zeta\left(\frac{3}{2}\right)(x\epsilon)^2\right] \\ &\quad +\left[-\frac{3\pi\sqrt{\epsilon}}{4}\coth(\pi\sqrt{\epsilon})+\frac{\pi^2\epsilon}{4}\operatorname{csch}^2(\pi\sqrt{\epsilon})\right]x^{-2}, \\ S(3, 1) &\sim \frac{15\sqrt{\pi}}{16}x^{-\frac{7}{2}}\left[\zeta\left(\frac{5}{2}\right)-\zeta\left(\frac{3}{2}\right)x\epsilon+\frac{1}{2}\zeta\left(\frac{1}{2}\right)(x\epsilon)^2\right] \\ &\quad +\left[\frac{5\pi\epsilon^{\frac{3}{2}}}{4}\coth(\pi\sqrt{\epsilon})-\frac{\pi^2\epsilon^2}{4}\operatorname{csch}^2(\pi\sqrt{\epsilon})\right]x^{-2}, \end{aligned} \quad (25)$$

where we consider $x\epsilon$ small and truncate the sum in (24) at the third term. This can be easily extended if necessary to higher orders.

We can now assess the accuracy of our approximation by comparing it with exact numerical results. We used (16) (with only the terms explicitly displayed) to determine ϵ , from which we calculate the fraction of particles in the ground state given by $N_{\text{gr}}/N = [e^{x(1+\epsilon)} - 1]^{-1}/N$ and the specific heat from (19) and (25). Fig. 1 shows N_{gr}/N for $N = 10^2$ and $N = 10^5$. Fig. 2 shows C/N for the same values of N . Had we also plotted the exact numerical results given by (19) and (20), the figures would look no different because the numerical plots would totally superimpose the analytical ones. For a better appreciation of the error involved in our analytical approximation see fig. 3, where we plot the error for N_{gr} .

Some comments are in order here. First, in all figures the scaling factor for the temperature is the infinite volume 3D critical temperature, T_c , given by $1/(2ma^2T_c) = x_c = [(\sqrt{\pi}\zeta(3/2))/(4N)]^{2/3}$. Given the volume of the three-sphere, $V = 2\pi^2a^3$, we can put T_c in terms of the particle density, ρ , as $T_c = (2\pi/m)(\rho/\zeta(3/2))^{2/3}$.

Secondly, we can clearly see from figs. 1 and 2 the approach to critical behaviour as N increases. In section IV we provide an analytical description for this phenomenon which explains the behaviour found here.

Finally, we consider the errors seen in fig. 3. We can see that these are very small for a wide range of x around the critical region. The peak of $\Delta N_{\text{gr}} = [N_{\text{gr}}(\text{ana}) - N_{\text{gr}}(\text{num})]/N_{\text{gr}}(\text{num})$ for $N = 10^5$ is a curious feature of our approximation. In a closer look, we can see this peak beginning to form already for $N = 10^2$. It increases with N . This feature can be misleading. While it is unquestionable that ΔN_{gr} increases with N in the critical region, the error in (16) for a given ϵ is actually diminishing. What happens is that as the thermodynamic limit is approached, we will have, for fixed $T < T_c$, $\epsilon \rightarrow -1$ and for fixed $T > T_c$, $\epsilon \rightarrow +\infty$, where T_c is the infinite volume critical temperature. This means that as N (and V) increases, ϵ will be an increasingly steeper function of temperature at T_c . This causes ϵ , given in (16) as an implicit function of N and x , to have an increasingly larger error in the critical region. The error is passed on to the ground state population. From fig. 3 we can also see that, excluding the already mentioned peak, the error increases as we get away from the centre (critical region). This is due to the series cut off at the third term in the relevant expressions. ($|x\epsilon|$ increases away from the centre.) By including more terms we can make the approximation work for an arbitrarily large range of x (or T). Similarly, neglecting more terms will cause the approximation to deteriorate faster as we get away from the centre. We do not plot the error in the specific heat, ΔC , because it does not show anything new. It behaves like ΔN_{gr} , except that the peak in the critical region is not manifest. This feature derives from C , as given in (19), being dominated by $S(3, 1)$, which in turn is dominated by the term in $x^{-7/2}$ in (25). This term does not depend on ϵ and so has a smooth behaviour.

B. The One-Dimensional Box

The energy eigenvalues for a non-relativistic spin-0 particle in a 1D box are

$$E_n = \frac{\pi^2}{2mL^2}n^2, \quad n = 1, 2, 3, \dots \quad (26)$$

where L is the length of the box. The levels are non-degenerate, ie, $d_n = 1$. After a change of variables,

$$x = \frac{\beta\pi^2}{2mL^2}, \quad \epsilon = -\frac{2mL^2}{\pi^2}\mu, \quad (27)$$

and using (1), we have

$$N = \sum_{n=1}^{+\infty} \left[e^{x(n^2+\epsilon)} - 1 \right]^{-1} = S(0,0). \quad (28)$$

Following the steps of the previous section yields

$$N \sim \frac{1}{2\pi i} \int_{\gamma} d\alpha \Gamma(\alpha) x^{-\alpha} \zeta(\alpha) g(\alpha, \epsilon). \quad (29)$$

For $\Gamma(\alpha)g(\alpha, \epsilon)$ the poles and respective residues are

$$\alpha = \frac{1}{2} - k, \quad k = 0, 1, 2, \dots \quad \text{res} = \frac{\sqrt{\pi}}{2} \frac{(-1)^k}{k!} \epsilon^k, \quad (30)$$

$$\alpha = -k, \quad k = 0, 1, 2, \dots \quad \text{res} = \frac{1}{2} \frac{(-1)^{k+1}}{k!} \epsilon^k, \quad (31)$$

resulting in

$$N \sim \frac{\sqrt{\pi}}{2} x^{-\frac{1}{2}} \sum_{k=0}^{+\infty} \frac{(-1)^k}{k!} \zeta\left(\frac{1}{2} - k\right) (\epsilon x)^k + \frac{1}{4} - \frac{1}{2} \sum_{k=1}^{+\infty} \frac{1}{(2k-1)!} \zeta(1-2k) (\epsilon x)^{2k-1} + x^{-1} g(1, \epsilon). \quad (32)$$

After truncating the sums and using (A3) we have,

$$N \sim \frac{\sqrt{\pi}}{2} x^{-\frac{1}{2}} \left[\zeta\left(\frac{1}{2}\right) + \frac{1}{4\pi} \zeta\left(\frac{3}{2}\right) \epsilon x - \frac{3}{32\pi^2} \zeta\left(\frac{5}{2}\right) (\epsilon x)^2 \right] + \frac{1}{4} - \frac{1}{24} \epsilon x + x^{-1} \left[\frac{\pi}{2\sqrt{\epsilon}} \coth(\pi\sqrt{\epsilon}) - \frac{1}{2\epsilon} \right]. \quad (33)$$

For the specific heat we can easily derive

$$C = x^2 \left[S(2,1) - \frac{S(1,1)^2}{S(0,1)} \right]. \quad (34)$$

We already have analytical expressions for $S(2,1)$ and $S(3,1)$. $S(0,1)$ is obtained in the same way, yielding

$$S(0,1) \sim \frac{\sqrt{\pi}}{2} x^{-\frac{1}{2}} \left[-\frac{1}{4\pi} \zeta\left(\frac{3}{2}\right) + \frac{3}{16\pi^2} \zeta\left(\frac{5}{2}\right) \epsilon x + \frac{15}{128\pi^3} \zeta\left(\frac{7}{2}\right) (\epsilon x)^2 \right] + x^{-2} \left[\frac{\pi^2}{4\epsilon} \text{csch}^2(\pi\sqrt{\epsilon}) + \frac{\pi}{4\epsilon^{3/2}} \coth(\pi\sqrt{\epsilon}) - \frac{1}{2\epsilon^2} \right], \quad (35)$$

where we use (A8).

The behaviour of the specific heat and ground state population is uninteresting in the 1D box case since all quantities are always smooth. It is well known [26,27] that for dimensions lower than three, there is no approach to BEC as $N \rightarrow +\infty$. For this reason, we do not include any plots here. The shapes of the curves can be easily extracted from the analytical expressions. In particular, in the large N limit we have

$$N = \frac{1}{x} \left[\frac{\pi}{2\sqrt{\epsilon}} \coth(\pi\sqrt{\epsilon}) - \frac{1}{2\epsilon} \right]. \quad (36)$$

As to the error contained in the approximation, it is smaller in this case than in S^3 . So that the reader will have an idea of just how small it is, we will restrict ourselves to saying that for $N = 10^3$, $\Delta N_{\text{gr}} = [N_{\text{gr}}(\text{ana}) - N_{\text{gr}}(\text{num})]/N_{\text{gr}}(\text{num})$ only becomes larger than 10^{-5} for $x^{-1} = 10^5$. Furthermore, the larger the value of N , the smaller the error for fixed x or fixed x/x_0 , where x_0 is defined as $\epsilon(x_0) = 0$, in analogy to x_0 in 3D (see section IV). This happens because, unlike in the Einstein universe, ϵ is bounded for any fixed x (or x/x_0).

C. The 1D Box $\otimes \mathbb{R}^2$

The 1Dbox $\otimes\mathbb{R}^2$ can be defined in several ways. One of them is to consider a 3D rectangular box with sides of length L_1 , L_2 and L_3 and then keep L_1 fixed and take $L_2 = cL_3 \rightarrow +\infty$ with $c \in \mathbb{R}^+$ being a constant. In a sense this is the most natural procedure and we shall adopt it.

The energy eigenvalues of a non-relativistic particle in such a box are

$$E_n = \frac{\pi^2}{2m} \left[\left(\frac{n_1}{L_1} \right)^2 + \left(\frac{n_2}{L_2} \right)^2 + \left(\frac{n_3}{L_3} \right)^2 \right], \quad n_i = 1, 2, 3 \dots \quad (37)$$

Using (1) and after some manipulation we have

$$N = \sum_{k=1}^{+\infty} \sum_{n_1=1}^{+\infty} e^{-k\beta \left(\frac{\pi^2 n_1^2}{2mL_1^2} - \mu \right)} \sum_{n_2=1}^{+\infty} e^{-k\beta \frac{\pi^2 n_2^2}{2mL_2^2}} \sum_{n_3=1}^{+\infty} e^{-k\beta \frac{\pi^2 n_3^2}{2mL_3^2}}. \quad (38)$$

Now, in the limit $L_2, L_3 \rightarrow +\infty$ the sums in n_2 and n_3 can be replaced by integrals, yielding

$$\sum_{n_i=1}^{+\infty} e^{-k\beta \frac{\pi^2 n_i^2}{2mL_i^2}} = \sqrt{\frac{m}{2\pi k\beta}} L_i, \quad i = 2, 3, \quad (39)$$

as the leading terms. Inserting (39) in (38) and making the change of variables

$$x = \frac{\beta\pi^2}{2mL_1^2}, \quad \epsilon = -\frac{2mL_1^2}{\pi^2}\mu, \quad (40)$$

we obtain

$$N = \frac{L_2 L_3}{L_1^2} \frac{\pi}{4} x^{-1} S(0, -1) \quad (41)$$

Since the volume is infinite when L_2 and L_3 are infinite, the number of particles is also infinite. It is then convenient to define a new quantity, η , which will be useful later, as

$$\eta = \frac{\pi}{4} x^{-1} S(0, -1). \quad (42)$$

η is the number of particles in a cube with sides of length L_1 . We prefer using η instead of the particle density, as η is the mathematical analogue of N in the previous cases.

Similarly to the previous sections, we obtain an analytical approximation for $S(0, -1)$ by applying a Mellin-Barnes transform, yielding

$$S(0, -1) \sim \frac{1}{2\pi i} \int_{\gamma} d\alpha \Gamma(\alpha) x^{-\alpha} \zeta(\alpha + 1) g(\alpha, \epsilon). \quad (43)$$

Extracting the residues of the poles of the integrand is a little trickier in this case, because the pole at $\alpha = 0$ is double. This pole comes from $\Gamma(\alpha)\zeta(\alpha + 1)$ since $g(\alpha, \epsilon)$ is finite at $\alpha = 0$. Furthermore,

$$\Gamma(\alpha)\zeta(\alpha + 1) = \frac{1}{\alpha^2} + \mathcal{O}(\alpha^0). \quad (44)$$

There is no term in α^{-1} (this is easily derived from Gamma and Riemann zeta function results in [25]). To get the residue we need the derivatives of $x^{-\alpha}$ and $g(\alpha, \epsilon)$ with respect to α . It will be

$$\text{res} = \left. \frac{\partial}{\partial \alpha} \right|_{\alpha=0} g(\alpha, \epsilon) - g(0, \epsilon) \ln x. \quad (45)$$

Using (A9) we have for this pole

$$\text{res} = \frac{1}{2} \ln x - \ln \frac{2 \sinh(\pi \sqrt{\epsilon})}{\sqrt{\epsilon}}. \quad (46)$$

The residues of the other poles are obtained in the manner of the previous sections, yielding

$$S(0, -1) \sim \frac{\sqrt{\pi}}{2} x^{-\frac{1}{2}} \left[\zeta\left(\frac{3}{2}\right) - \zeta\left(\frac{1}{2}\right) \epsilon x - \frac{1}{8\pi} \zeta\left(\frac{3}{2}\right) (\epsilon x)^2 \right] - \frac{1}{4} \epsilon x + \frac{1}{2} \ln x - \ln \frac{2 \sinh(\pi \sqrt{\epsilon})}{\sqrt{\epsilon}} . \quad (47)$$

Like in the Einstein universe, this can be seen to approach the continuum approximation when $x \rightarrow 0$.

Now, for the specific heat, the analogue of expressions (19) and (34) here is

$$C = \frac{\pi L_2 L_3}{4 L_1^2} \left[x S(2, 0) + 2 S(1, -1) + \frac{2}{x} S(0, -2) - x \frac{[\frac{1}{x} S(0, -1) + S(1, 0)]^2}{S(0, 0)} \right] . \quad (48)$$

In the course of this article we already obtained analytical expressions for $S(0, 0)$ (1D box number of particles), $S(1, 0)$ (S^3 number of particles) and $S(0, -1)$ (1D box $\otimes \mathbb{R}^2$ number of particles). We now need expressions for $S(2, 0)$, $S(1, -1)$ and $S(0, -2)$.

For $S(0, -2)$ the procedure is identical to the one used to get $S(0, -1)$. Now the double pole is at $\alpha = -1$ and its residue is

$$\text{res} = \frac{1}{2} x \epsilon - \frac{1}{2} x \epsilon \ln x - \frac{\partial}{\partial \alpha} \Big|_{\alpha=-1} g(\alpha, \epsilon) . \quad (49)$$

The expression for $\partial g / \partial \alpha$ at $\alpha = -1$, given by (A11), is not as simple as we would have liked, as it includes an infinite sum. However, that turns out not to be limiting when it comes to studying analytically the critical region, as we will see. Using (A11) we obtain

$$S(0, -2) \sim \frac{\sqrt{\pi}}{2} x^{-\frac{1}{2}} \left[\zeta\left(\frac{5}{2}\right) - \zeta\left(\frac{3}{2}\right) \epsilon x + \frac{1}{2} \zeta\left(\frac{1}{2}\right) (\epsilon x)^2 \right] - \frac{\pi^2}{12} + \frac{1}{2} \epsilon x - \frac{1}{2} \epsilon x \ln x + \frac{\zeta(3)}{2\pi^2} x + \epsilon x \ln \frac{2 \sinh(\pi \sqrt{\epsilon})}{\sqrt{\epsilon}} - x \sum_{k=0}^{+\infty} \frac{(2\pi)^{2k} B_{2k}}{(2+2k)(2k)!} \epsilon^{k+1} , \quad (50)$$

for $|\epsilon| < 1$. B_{2k} are the Bernoulli numbers.

$S(2, 0)$ is simpler. All the poles are simple and it does not present any additional complications in relation to the sums studied in the previous sections. The result is

$$S(2, 0) \sim \frac{3\sqrt{\pi}}{8} x^{-\frac{5}{2}} \left[\zeta\left(\frac{5}{2}\right) - \zeta\left(\frac{3}{2}\right) \epsilon x + \frac{1}{2} \zeta\left(\frac{1}{2}\right) (\epsilon x)^2 \right] + \frac{\pi \epsilon^{\frac{3}{2}}}{2} \coth(\pi \sqrt{\epsilon}) x^{-1} . \quad (51)$$

In the case of $S(1, -1)$, there are not any double poles either. Still, it deserves a little attention. Proceeding as before, we have

$$S(1, -1) \sim \frac{1}{2\pi i} \int_{\gamma} d\alpha \Gamma(\alpha) x^{-\alpha} \zeta(\alpha + 1) f_1(\alpha, \epsilon) . \quad (52)$$

The pole deriving from the zeta function is at $\alpha = 0$. Despite the fact that $\Gamma(\alpha)$ has a pole at $\alpha = 0$, the function $\Gamma(\alpha) f_1(\alpha, \epsilon)$ can be analytically continued there. The reason is that $f_1(0, \epsilon) = 0$, cancelling the $\Gamma(\alpha)$ pole. The residue of the pole of $\Gamma(\alpha)$ at $\alpha = 0$ is 1. We have then,

$$\begin{aligned} \lim_{\alpha \rightarrow 0} \Gamma(\alpha) f_1(\alpha, \epsilon) &= \frac{\partial}{\partial \alpha} \Big|_{\alpha=0} f_1(\alpha, \epsilon) \\ &= -\frac{\zeta(3)}{2\pi^2} + \sum_{k=0}^{+\infty} \frac{(2\pi)^{2k} B_{2k}}{(2+2k)(2k)!} \epsilon^{k+1} . \end{aligned} \quad (53)$$

The last equality is valid for $|\epsilon| < 1$ (see appendix). Finally we arrive at

$$S(1, -1) \sim \frac{\sqrt{\pi}}{4} x^{-\frac{3}{2}} \left[\zeta\left(\frac{5}{2}\right) - \zeta\left(\frac{3}{2}\right) \epsilon x + \frac{1}{2} \zeta\left(\frac{1}{2}\right) (\epsilon x)^2 \right] - \frac{\zeta(3)}{2\pi^2} + \sum_{k=0}^{+\infty} \frac{(2\pi)^{2k} B_{2k}}{(2+2k)(2k)!} \epsilon^{k+1}. \quad (54)$$

We can now obtain ϵ from (42) and (47) and use it to calculate η_1 , which is the analog of N_{gr} in the Einstein universe:

$$\begin{aligned} \eta_1 &= \frac{\pi}{4} x^{-1} \sum_{k=1}^{+\infty} \frac{e^{-kx(1+\epsilon)}}{k} \\ &= -\frac{\pi}{4} x^{-1} \ln \left[1 - e^{-x(1+\epsilon)} \right]. \end{aligned} \quad (55)$$

η_1 is obtained from (42) by considering only the $n = 1$ part of the sum $S(0, -1)$. It is not the ground state population. The fraction of particles in the ground state has no interest to us, since N_{gr} is finite (expression (38) with $n_1 = n_2 = n_3 = 1$) and so, when L_2 and L_3 are infinite, $N_{\text{gr}}/N = 0$. In contrast, η_1/η is the relevant quantity here. It has a similar behaviour to N_{gr}/N in the Einstein universe.

Making $L_1 \rightarrow +\infty$ in our system is a way of obtaining the infinite three dimensional space. And yet, we have $N_{\text{gr}}/N \rightarrow 0$ for all temperatures when $L_1 \rightarrow +\infty$. This shows that the statement that in infinite 3D, N_{gr}/N is finite below a certain critical temperature is incorrect. We must specify how we make our 3D space infinite. In the statement above, we should replace “infinite 3D” by, for example, “rectangular box of sides L_1, L_2 and L_3 with $L_1, L_2, L_3 \rightarrow +\infty$ while L_1/L_2 and L_2/L_3 are kept constant”. One might rightly argue that the space we study here is not a box, since L_2 and L_3 are taken to infinity from the start. However, it is possible to show that we can have a 3D box for which $N_{\text{gr}}/N \rightarrow 0$ for all temperatures when $L_1, L_2, L_3 \rightarrow +\infty$ if the lengths of the sides obey particular relations between them.

In fig. 1 we plot η_1/η for $\eta = 10^2$ and $\eta = 10^5$. Only the analytical curves are plotted. Like in the Einstein universe, if the numerical curves were plotted, they would totally superimpose the analytical ones. In fig. 2 we plot C/N obtained from (48), (16), (33), (47), (50), (51) and (54). The point where the curves change the pattern in fig. 2 is where ϵ becomes larger than 1, rendering (50) and (54) not valid anymore. Both the analytical curves, obtained from the expressions mentioned above, and the exact numerical ones, obtained from (42), (48) and (20), are plotted. In the range where the analytical approximation applies ($\epsilon < 1$), they are totally superimposed, looking like one curve only. For $\epsilon > 1$ only the numerical curves are plotted. In both figs. 1 and 2, the curves are similar to those of the Einstein universe. In the large N limit they tend to be the same, specifically, the standard text book curves of the infinite box. This can be seen analytically by taking $x \rightarrow 0$ in our approximations.

In fig. 3 we plot the error $\Delta\eta_1 = [\eta_1(\text{ana}) - \eta_1(\text{num})]/\eta_1(\text{num})$. All the error analysis done for the Einstein universe applies here also.

III. A NOTE ON THE MELLIN-BARNES TRANSFORM

As we mentioned already, when going from expression (7) to (8) we neglect the contribution from the added part of the contour to the integral. Although this contribution is vanishing for example for expression (6) [28], it is not necessarily so for (7). It is at most very small, as we could see when comparing the analytical results to the exact numerical ones. However, the peak in fig. 3 clearly indicates that it is not vanishing. It is smaller and smaller as N gets larger at fixed density (and hence, x gets smaller), but if it was null the peak would not be there at all.

It is possible, if this contribution is not negligible, to have a situation in which the Mellin-Barnes transform technique will give totally wrong results. In our investigations, we came across such an example. When applying the transform to (4) instead of applying it to the whole exponential, we could have split the exponential in two, $e^{-kx(n^2-1)}e^{-kx(\epsilon+1)}$, take the ground state out [29], and apply the transform to the first part only, in the manner of [16]. This yields

$$N = N_{\text{gr}} + \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} d\alpha \Gamma(\alpha) x^{-\alpha} \text{Li}_{\alpha}(e^{-x(\epsilon+1)}) f(\alpha), \quad (56)$$

where $f(\alpha)$ and the polylogarithm $\text{Li}_{\alpha}(a)$ are defined like

$$f(\alpha) = \sum_{n=1}^{+\infty} (n+1)^2 [n(n+2)]^{-\alpha},$$

$$\text{Li}_\alpha(a) = \sum_{k=1}^{+\infty} \frac{a^k}{k^\alpha} . \quad (57)$$

Closing the contour to the left and proceeding as in the previous section, we obtain

$$N \sim \frac{\sqrt{\pi}}{4} x^{-\frac{3}{2}} \sum_{k=0}^{+\infty} \frac{1}{k!} \text{Li}_{\frac{3}{2}-k}(e^{-x(\epsilon+1)}) x^k . \quad (58)$$

For $\epsilon < 0$, this expression is completely wrong. This can be seen by noting that

$$\sum_{k=0}^{+\infty} \frac{1}{k!} \text{Li}_{\frac{3}{2}-k}(e^{-x(\epsilon+1)}) x^k = \text{Li}_{\frac{3}{2}}(e^{-x\epsilon}) . \quad (59)$$

Therefore, our expression is divergent for $\epsilon < 0$. Curiously, this is exactly the same as the continuum approximation (also called “bulk”), ie, directly replacing the sum in n in (4) by an integral.

This way of applying the Mellin-Barnes transform is incorrect not only in the Einstein universe case. It is not difficult to show that if the energy levels are of the form $E_n - E_0 = a(n+b)^i + c$, $a \in \mathbb{R}^+$, $i \in \mathbb{R}^+$, $b \in \mathbb{R} \setminus \mathbb{Z}^-$, $c \in \mathbb{R}$ and the levels degeneracy, d_n , is any polynomial (and this includes all the cases in this article), then the situation will be identical. In principle, there is nothing preventing this from generalizing to more cases still. Thus, one must be very careful about how to apply the Mellin-Barnes transform.

IV. CRITICAL REGION EXPANSIONS

Expressions (16) and (47) give us ϵ implicitly and from here we have all thermodynamic quantities. However, it would be much more convenient to have ϵ given explicitly. That is what we will aim for in this section, concentrating on the critical region. This is similar to what was done in [18] in the context of a gas subject to a magnetic field. However, an important difference is that we do not perform our expansions around x_c . Instead, we use x_0 (defined below), making the problem considerably easier to treat. This will still give us the critical behaviour since when $N \rightarrow +\infty$ we have $x_0 \rightarrow x_c$. We then use the results obtained on the specific heat.

A. Einstein Universe

Define x_0 as being the value of x at which $\epsilon = 0$. From (16), we have

$$N = \frac{\sqrt{\pi}}{4} \zeta\left(\frac{3}{2}\right) x_0^{-\frac{3}{2}} - \frac{1}{2} x_0^{-1} . \quad (60)$$

Still in (16), change variables from x to $\delta = (x/x_0)^{1/2} - 1$. The Taylor expansion of ϵ as a function of δ (for fixed N) is

$$\epsilon = a_1 \delta + a_2 \delta^2 + \dots , \quad (61)$$

where the a_i s are, for the moment, unknown. There is no independent term since by construction $\epsilon(0) = 0$. We now expand $\coth(\pi\sqrt{\epsilon})$ around $\sqrt{\epsilon} = 0$ and insert (60) and (61) in (16). Solving for each power of δ finally yields

$$a_1 = -x_0^{-\frac{1}{2}} \frac{\frac{3\sqrt{\pi}}{4} \zeta\left(\frac{3}{2}\right) - x_0^{\frac{1}{2}}}{\frac{\pi^2}{6} + \frac{\sqrt{\pi}}{4} \zeta\left(\frac{1}{2}\right) x_0} . \quad (62)$$

For large N , $x_0 \ll 1$ and we have to highest order in x_0^{-1} ,

$$a_1 \simeq -\frac{9\zeta\left(\frac{3}{2}\right)}{2\pi^{\frac{3}{2}}} x_0^{-\frac{1}{2}} . \quad (63)$$

The result for a_2 is quite long and there is no point in displaying it in full. For $x_0 \ll 1$ it becomes

$$a_2 \simeq \frac{27\zeta\left(\frac{3}{2}\right)^2}{20\pi}x_0^{-1}. \quad (64)$$

Expressions (61), (63) and (64) give us ϵ explicitly in a neighbourhood of x_0 .

We can now find an expansion for the specific heat by inserting (61) in (25), obtaining

$$\begin{aligned} \frac{C}{N} &= \frac{15\zeta\left(\frac{5}{2}\right)}{4\zeta\left(\frac{3}{2}\right)} + \left[\frac{81\zeta\left(\frac{3}{2}\right)^2}{40\pi} - \frac{45\zeta\left(\frac{5}{2}\right)}{4\zeta\left(\frac{3}{2}\right)} \right] \delta - \frac{729\zeta\left(\frac{3}{2}\right)^3}{1400\sqrt{\pi}}x_0^{-\frac{1}{2}}\delta^2 + \dots \\ &\simeq 1.93 - 1.38\delta - 5.24x_0^{-\frac{1}{2}}\delta^2 + \dots \end{aligned} \quad (65)$$

The coefficients of each term are given only to highest order in x_0^{-1} . The first term is the familiar specific heat maximum of the 3D infinite space. The other terms are dependent on which kind of 3D space we have, ie, they are specific to the Einstein universe. From here we immediately get the first and second derivatives of C with respect to δ at $\delta = 0$ and hence, with respect to T . The first derivative is a constant, but the second one goes to infinity as $N \rightarrow +\infty$ ($x_0 \rightarrow 0$). This is not surprising, it is the famous specific heat peak getting sharper. It does not mean that T_0 is the same as the critical temperature T_c . What we have is $\lim_{N \rightarrow +\infty} T_0 = T_c$ (which is already evident from (60)). However, if we define T_a as the temperature at which $\epsilon = a \neq 0$, we will still have $\lim_{N \rightarrow +\infty} T_a = T_c$ because the function $\epsilon(T)$ tends to become infinitely steep at T_c . $\epsilon = 0$ is, nevertheless, a privileged choice as it provides the simplest treatment for the problem.

For completeness, we also present C/N at T_0 expanded around $x_0 = 0$ ($N = +\infty$):

$$\begin{aligned} \frac{C}{N}(x_0) &= \frac{15\zeta\left(\frac{5}{2}\right)}{4\zeta\left(\frac{3}{2}\right)} + \left[\frac{15\zeta\left(\frac{5}{2}\right)}{2\sqrt{\pi}\zeta\left(\frac{3}{2}\right)^2} - \frac{27\zeta\left(\frac{3}{2}\right)}{8\pi^{\frac{3}{2}}} \right] x_0^{\frac{1}{2}} + \dots \\ &\simeq 1.93 - 0.75x_0^{\frac{1}{2}} + \dots \end{aligned} \quad (66)$$

B. 1D Box $\otimes \mathbb{R}^2$

For the 1Dbox $\otimes\mathbb{R}^2$ the process is exactly the same. In this case, x_0 is still the value of x at which $\epsilon = 0$ and is given by (42) and (47) as

$$\eta = \frac{\sqrt{\pi}}{4}\zeta\left(\frac{3}{2}\right)x_0^{-\frac{3}{2}} + \frac{1}{4}x_0^{-1}\ln(x_0) - \frac{1}{2}x_0^{-1}\ln(2\pi). \quad (67)$$

Again, we have that $\lim_{N \rightarrow +\infty} x_0 = x_c$.

The result for ϵ is

$$\epsilon = -\frac{9\zeta\left(\frac{3}{2}\right)}{\pi^{\frac{3}{2}}}x_0^{-\frac{1}{2}}\delta + \frac{27\zeta\left(\frac{3}{2}\right)^2}{10\pi}x_0^{-1}\delta^2 + \dots \quad (68)$$

where $\delta = \sqrt{x/x_0} - 1$. The coefficients are only displayed to highest order in x_0^{-1} .

For the specific heat, inserting (68) in (16), (33), (47), (50), (51) and (54) we have

$$\begin{aligned} \frac{C}{N} &= \frac{15\zeta\left(\frac{5}{2}\right)}{4\zeta\left(\frac{3}{2}\right)} + \left[\frac{81\zeta\left(\frac{3}{2}\right)^2}{20\pi} - \frac{45\zeta\left(\frac{5}{2}\right)}{4\zeta\left(\frac{3}{2}\right)} \right] \delta - \frac{243\zeta\left(\frac{3}{2}\right)^3}{1400\sqrt{\pi}}x_0^{-\frac{1}{2}}\delta^2 + \dots \\ &\simeq 1.93 + 3.02\delta - 1.75x_0^{-\frac{1}{2}}\delta^2 + \dots \end{aligned} \quad (69)$$

Expressions (50) and (54) are valid only for $|\epsilon| < 1$. Since we are expanding around $\epsilon = 0$, this is not important, but it means that (69) is valid only for $\delta > \delta'$, where δ' is a negative number satisfying $\epsilon(\delta') = 1$. Comparing (69) with (65) we see that the qualitative behaviour is very similar except that now the second term has a positive coefficient, meaning that T_0 is to the right of the specific heat maximum, as opposed to the previous case. The second derivative still becomes infinitely negative as $x_0 \rightarrow 0$. Once more, this corresponds to the sharpening of the specific heat peak.

The equivalent of (66) here is

$$\begin{aligned} \frac{C}{N}(x_0) &= \frac{15\zeta\left(\frac{5}{2}\right)}{4\zeta\left(\frac{3}{2}\right)} - \frac{15\zeta\left(\frac{5}{2}\right)}{4\sqrt{\pi}\zeta\left(\frac{3}{2}\right)^2}x_0^{\frac{1}{2}}\ln\frac{x_0}{(2\pi)^2} - \left[\frac{\pi^{\frac{3}{2}}}{3\zeta\left(\frac{3}{2}\right)} + \frac{27\zeta\left(\frac{3}{2}\right)}{4\pi^{\frac{3}{2}}} \right] x_0^{\frac{1}{2}} + \dots \\ &\simeq 1.93 - 0.42x_0^{\frac{1}{2}}\ln x_0 - 2.35x_0^{\frac{1}{2}} + \dots \end{aligned} \quad (70)$$

V. CONCLUSION

We have provided a general method to treat some thermodynamic problems by invoking the Mellin-Barnes transform. In principle, it can be used to treat any problem that presents the same kind of sums as long as the quantity equivalent to ϵx does not become very large in the range of interest. Although we have used it in systems with quadratic energy levels only, it should be possible to use it in other systems as well. This method is inspired by the one in [16] with some important modifications. Moreover, we have shown that the method presented in [16] has limitations. In particular, it is not applicable to the spaces studied here and generally to systems with energy eigenvalues of some specific forms. Perhaps it would be interesting to look at this from the point of view of the density of states method which was shown in [16] to be linked to the Mellin-Barnes technique.

As already mentioned, the results obtained in section II for the number of particles in the Einstein universe could have been obtained using the Poisson summation formula as in [8]. It is likely that we could have used this technique to get all the results of section II. It would be interesting to establish a deeper connection between the two techniques.

We have also provided an analytical description of the appearance of the sharp specific heat peak and hence, the onset of BEC. As far as we know, this has not been done before. It was a natural consequence of obtaining an analytical result for C , expanded around the critical region, in terms of the temperature only and not the chemical potential, which in turn, derived from having the chemical potential as an explicit function of the temperature, in a similar fashion to [18]. Our particular choice of temperature around which to perform the expansions simplified the calculations and therefore it was possible to obtain the results for the specific heat.

As a side issue, from our treatment of the $1\text{Dbox} \otimes \mathbb{R}^2$ arose the conceptual problem of what one means by infinite three-dimensional space. We pointed out that the way this limit is taken must be specified before we can make certain statements about the system. In particular, the system treated is an example of the onset of BEC at a finite and not vanishing critical temperature while the fraction of particles in the ground state (N_{gr}/N) remains zero at all temperatures. Still, there will be a quantity that takes the standard role of N_{gr}/N . In our case, it is the fraction of particles that has minimum frequency ($n_1 = 1$) in the L_1 direction.

ACKNOWLEDGMENTS

J.N. acknowledges financial support by the Portuguese Foundation for Science and Technology under grant Prax-isXXI/BD/5660/95, which extended through part of the period encompassed by this work.

APPENDIX:

Consider the function defined, for $i \in \mathbb{N}^0$, as

$$f_i(\alpha, \epsilon) = \sum_{n=1}^{+\infty} n^{2i} (n^2 + \epsilon)^{-\alpha} \quad \text{for } \Re(\alpha) > i + \frac{1}{2}. \quad (\text{A1})$$

When $\Re(\alpha) \leq i + 1/2$, we will define $f_i(\alpha, \epsilon)$ by analytic continuation as described below. In this appendix we want to obtain f_i and $\partial f_i / \partial \alpha$ for some special values of i and α .

Define $g \equiv f_0$. The analytical continuation of this function can be achieved by noting that

$$\frac{\partial}{\partial \epsilon} g(\alpha, \epsilon) = -\alpha g(\alpha + 1, \epsilon). \quad (\text{A2})$$

For $\Re(\alpha) > 1/2$, this follows immediately from (A1). In order to have an analytical $\partial g / \partial \epsilon$, this equality must extend to the complex plane. $g(1, \epsilon)$ is given by

$$\begin{aligned} g(1, \epsilon) &= \sum_{n=1}^{+\infty} (n^2 + \epsilon)^{-1} \\ &= \frac{\pi}{2\sqrt{\epsilon}} \coth(\pi\sqrt{\epsilon}) - \frac{1}{2\epsilon} \end{aligned} \quad (\text{A3})$$

or

$$g(1, \epsilon) = -\frac{\pi}{2\sqrt{-\epsilon}} \cot(\pi\sqrt{-\epsilon}) - \frac{1}{2\epsilon} . \quad (\text{A4})$$

Both expressions (A3) and (A4) are valid for $\epsilon > 0$ and $\epsilon < 0$, but the first one is more convenient for $\epsilon > 0$ and the second for $\epsilon < 0$. $g(0, \epsilon)$ is obtained by integrating (A2) with $\alpha = 0$, yielding

$$g(0, \epsilon) = g(0, 0) . \quad (\text{A5})$$

We know that $g(\alpha, 0) \equiv \zeta(2\alpha)$. Thus,

$$g(0, \epsilon) = \zeta(0) = -\frac{1}{2} . \quad (\text{A6})$$

$g(-1, \epsilon)$ is obtained by using (A2) with $\alpha = -1$ in conjunction with (A6), resulting in

$$g(-1, \epsilon) = -\frac{1}{2}\epsilon . \quad (\text{A7})$$

We also need $g(2, \epsilon)$. This comes directly from (A2). Putting $\alpha = 1$, we have

$$g(2, \epsilon) = \frac{\pi^2}{4\epsilon} \text{csch}^2(\pi\sqrt{\epsilon}) + \frac{\pi}{4\epsilon^{3/2}} \coth(\pi\sqrt{\epsilon}) - \frac{1}{2\epsilon^2} . \quad (\text{A8})$$

Now, we want the derivatives $\partial g/\partial\alpha$ at $\alpha = 0$ and $\alpha = -1$. Inserting the Taylor series of $g(\alpha, \epsilon)$ and $g(1 + \alpha, \epsilon)$ in (A2) and integrating, yields

$$\begin{aligned} \left. \frac{\partial}{\partial\alpha} \right|_{\alpha=0} g(\alpha, \epsilon) &= 2\zeta'(0) - \int_0^\epsilon g(1, \epsilon) d\epsilon \\ &= -\ln \frac{2 \sinh(\pi\sqrt{\epsilon})}{\sqrt{\epsilon}} , \end{aligned} \quad (\text{A9})$$

where the prime in $\zeta'(0)$ means derivative and we used $\zeta'(0) = -(\ln 2\pi)/2$ and (A3). Following the same procedure for $\partial g/\partial\alpha$ at $\alpha = -1$, we obtain

$$\left. \frac{\partial}{\partial\alpha} \right|_{\alpha=-1} g(\alpha, \epsilon) = 2\zeta'(-2) + \frac{1}{2}\epsilon - \int_0^\epsilon \ln \frac{2 \sinh(\pi\sqrt{\epsilon})}{\sqrt{\epsilon}} d\epsilon . \quad (\text{A10})$$

There is no simple form for the integral in (A10). However, in the case of $|\epsilon| < 1$, it can be expressed in terms of an infinite sum (see eg [25]), which is more convenient than just leaving it as it is. After getting $\zeta'(-2) = -\zeta(3)/(2\pi^2)$ from expressions in [25], we finally have

$$\left. \frac{\partial}{\partial\alpha} \right|_{\alpha=-1} g(\alpha, \epsilon) = -\frac{\zeta(3)}{2\pi^2} - \epsilon \ln \frac{2 \sinh(\pi\sqrt{\epsilon})}{\sqrt{\epsilon}} + \sum_{k=0}^{+\infty} \frac{(2\pi)^{2k} B_{2k}}{(2+2k)(2k)!} \epsilon^{k+1} \quad (\text{A11})$$

(valid for $|\epsilon| < 1$), where B_{2k} are the Bernoulli numbers.

The results for $f_i(\alpha, \epsilon)$, $i \in \mathbb{N}$ are now easy to obtain. These functions can be expressed in terms of $g(\alpha, \epsilon)$, as can be seen for $\Re(\alpha) > i + 1/2$ in (A1). For $f_1(\alpha, \epsilon)$ we have

$$f_1(\alpha, \epsilon) = g(\alpha - 1, \epsilon) - \epsilon g(\alpha, \epsilon) \quad (\text{A12})$$

This equality must hold also for $\Re(\alpha) \leq i + 1/2$, so that $f_1(\alpha, \epsilon)$ is analytical. The extension to $i \neq 1$ does not present any additional difficulties. The special values we will need are

$$f_1(0, \epsilon) = 0 , \quad (\text{A13})$$

$$f_1(1, \epsilon) = -\frac{\pi\sqrt{\epsilon}}{2} \coth(\pi\sqrt{\epsilon}) , \quad (\text{A14})$$

$$f_1(2, \epsilon) = \frac{\pi}{4\sqrt{\epsilon}} \coth(\pi\sqrt{\epsilon}) - \frac{\pi^2}{4} \text{csch}^2(\pi\sqrt{\epsilon}) , \quad (\text{A15})$$

$$f_2(1, \epsilon) = \frac{\pi\epsilon^{\frac{3}{2}}}{2} \coth(\pi\sqrt{\epsilon}) , \quad (\text{A16})$$

$$f_2(2, \epsilon) = -\frac{3\pi\sqrt{\epsilon}}{4} \coth(\pi\sqrt{\epsilon}) + \frac{\pi^2\epsilon}{4} \operatorname{csch}^2(\pi\sqrt{\epsilon}) , \quad (\text{A17})$$

$$f_3(2, \epsilon) = \frac{5\pi\epsilon^{\frac{3}{2}}}{4} \coth(\pi\sqrt{\epsilon}) - \frac{\pi^2\epsilon^2}{4} \operatorname{csch}^2(\pi\sqrt{\epsilon}) , \quad (\text{A18})$$

$$\left. \frac{\partial}{\partial \alpha} \right|_{\alpha=0} f_1(\alpha, \epsilon) = -\frac{\zeta(3)}{2\pi^2} + \sum_{k=0}^{+\infty} \frac{(2\pi)^{2k} B_{2k}}{(2+2k)(2k)!} \epsilon^{k+1} . \quad (\text{A19})$$

Expression (A19) is valid only for $|\epsilon| < 1$.

- [1] Y. Castin, in “*Coherent Matter Waves*”, *Lecture Notes of Les Houches 1999 Summer School, Session LXXII*, edited by R. Kaiser, C. Westbrook and F. David (EDP Sciences and Springer-Verlag, 2001); F. Dalfovo, S. Giorgini, L. P. Pitaevskii and S. Stringari *Rev. Mod. Phys.* **71**, 463 (1999).
- [2] W. Ketterle and E. Cornell, in “*Coherent Matter Waves*”, *Lecture Notes of Les Houches 1999 Summer School, Session LXXII*, edited by R. Kaiser, C. Westbrook and F. David (EDP Sciences and Springer-Verlag, 2001).
- [3] S. Grossmann and M. Holthaus, *Phys. Lett. A* **208**, 188 (1995).
- [4] W. Ketterle and N. J. van Druten, *Phys. Rev. A* **54**, 656 (1996).
- [5] K. Kirsten and D. J. Toms, *Phys. Lett. A* **222**, 148 (1996).
- [6] K. Kirsten and D. J. Toms, *Phys. Rev. A* **54**, 4188 (1996).
- [7] H. Haugerud, T. Haugset and F. Ravndal, *Phys. Lett. A* **225**, 18 (1997); T. Haugset, H. Haugerud and J. O. Andersen, *Phys. Rev. A* **55**, 2922 (1997).
- [8] M. B. Altaie, *J. Phys. A: Math. Gen.* **11**, 1603 (1978).
- [9] S. Singh and R. K. Pathria, *J. Phys. A: Math. Gen.* **17**, 2983 (1984).
- [10] M. Trucks, *Phys. Lett. B* **445**, 117 (1998).
- [11] M. B. Altaie and E. Malkawi, *J. Phys. A: Math. Gen.* **33**, 7093 (2000).
- [12] L. Parker and Y. Zhang, *Phys. Rev. D* **44**, 2421 (1991).
- [13] D. J. Toms, *Phys. Rev. Lett.* **69**, 1152 (1992); D. J. Toms, *Phys. Rev. D* **47**, 2483 (1993).
- [14] K. Kirsten and D. J. Toms, *Phys. Rev. D* **51**, 6886 (1995).
- [15] J. D. Smith and D. J. Toms, *Phys. Rev. D* **53**, 5771 (1996).
- [16] K. Kirsten and D. J. Toms, *Phys. Lett. A* **243**, 137 (1998); K. Kirsten and D. J. Toms, *Phys. Rev. E* **59**, 158 (1999).
- [17] F. Brosens, J. T. Devreese and L. F. Lemmens, *Solid State Comm.* **100**, 123 (1996).
- [18] G. B. Standen and D. J. Toms, *Phys. Rev. E* **60**, 5275 (1999).
- [19] M. Holthaus, E. Kalinowski and K. Kirsten, *Annals of Physics* **270**, 198 (1998).
- [20] H. D. Politzer, *Phys. Rev. A* **54**, 5048 (1996).
- [21] M. Gajda and K. Rzążewski, *Phys. Rev. Lett.* **78**, 2686 (1997).
- [22] M. Wilkens and C. Weiss, *J. Mod. Optics* **44**, 1801 (1997).
- [23] S. Grossmann and M. Holthaus, e-print cond-mat/9709045
- [24] P. Borrmann, J. Harting, O. Mulken and E. R. Hilf, *Phys. Rev. A* **60**, 1519 (1999).
- [25] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals Series and Products* (Academic Press, 1994).
- [26] R. M. May, *Phys. Rev.* **115**, 254 (1959).
- [27] K. Kirsten and D. J. Toms, *Phys. Lett. B* **368**, 119 (1996).
- [28] In fact, this is how you prove (6) (see [6]).
- [29] The reason we do the split like this and not $e^{-kxn^2}e^{-kx\epsilon}$ and take the ground state out is to allow the separation of the sums in k and n , with each of them remaining convergent.




FIG. 1. Analytical N_{gr}/N in the Einstein universe for $N = 10^2$ (continuous line) and $N = 10^5$ (dotted line) and η_1/η in the $1\text{Dbox} \otimes \mathbb{R}^2$ for $\eta = 10^2$ (dashed line) and $\eta = 10^5$ (dashed/dotted line) as functions of the rescaled temperature




FIG. 2. The specific heat per particle in the Einstein universe for $N = 10^2$ (analytical; dashed line) and $N = 10^5$ (analytical; dashed/dotted line) and in the $1D\text{box} \otimes \mathbb{R}^2$ for $\eta = 10^2$ (continuous -analytical- and then dashed -numerical- line) and $\eta = 10^5$ (dashed -analytical- and then dotted -numerical- line) - see text for details




FIG. 3. ΔN_{gr} in the Einstein universe for $N = 10^2$ (continuous line) and $N = 10^5$ (dotted line) and $\Delta \eta_1$ in the $1\text{Dbox} \otimes \mathbb{R}^2$ for $\eta = 10^2$ (dashed line) and $\eta = 10^5$ (dashed/dotted line) - see text for the definitions of ΔN_{gr} and $\Delta \eta_1$